



InterClinical Laboratories Pty Limited
 ABN 89 076 386 475
 PO Box 6474, Alexandria NSW 2015 Australia
 Ph: 02 9693 2888 Fax: 02 9693 1888
 Email: lab@interclinical.com.au


LABORATORY NO.: **XXXXX**

PROFILE NO.: **2** SAMPLE TYPE: **SCALP**


PATIENT: **XXXXX** AGE: **69** SEX: **M** METABOLIC TYPE: **FAST 4**

REQUESTED BY: **XXXXX** ACCOUNT NO.: **2216** DATE: **2021**


NUTRITIONAL ELEMENTS



TOXIC ELEMENTS




ADDITIONAL ELEMENTS



"<<": Below Calibration Limit; Value Given Is Calibration Limit
 "QNS": Sample Size Was Inadequate For Analysis.
 "N/A": Currently Not Available
 Ideal Levels And Interpretation Have Been Based On Hair Samples Obtained From The Mid-Parietal To The Occipital Region Of The Scalp.
 Laboratory Analysis Provided by Trace Elements, Inc. Dallas, Texas USA an H.H.S. Licensed Clinical Laboratory. No. 45 D0481787

2021
 CURRENT TEST RESULTS
 PREVIOUS TEST RESULTS

SIGNIFICANT RATIOS



TOXIC RATIOS



ADDITIONAL RATIOS

RATIO	CALCULATED VALUE		EXPECTED
	Current	Previous	
Ca/Sr	533.33		263/1
Cr/V	10.00		8/1
Cu/Mo	1600.00		356/1
Fe/Co	400.00		615/1
K/Co	2000.00		6350/1
K/Li	2000.00		6350/1
Mg/B	N/A		21/1
S/Cu	2628.13		2668/1
Se/Tl	600.00		370/1
Se/Sn	6.00		3.2/1
Zn/Sn	1400.00		624/1

LEVELS

All mineral levels are reported in milligrams percent (milligrams per one-hundred grams of hair). One milligram percent (mg%) is equal to ten parts per million (ppm).

NUTRITIONAL ELEMENTS

Extensively studied, the nutrient elements have been well defined and are considered essential for many biological functions in the human body. They play key roles in such metabolic processes as muscular activity, endocrine function, reproduction, skeletal integrity and overall development.

TOXIC ELEMENTS

The toxic elements or "heavy metals" are well-known for their interference upon normal biochemical function. They are commonly found in the environment and therefore are present to some degree, in all biological systems. However, these metals clearly pose a concern for toxicity when accumulation occurs to excess.

ADDITIONAL ELEMENTS

These elements are considered as possibly essential by the human body. Additional studies are being conducted to better define their requirements and amounts needed.

RATIOS

A calculated comparison of two elements to each other is called a ratio. To calculate a ratio value, the first mineral level is divided by the second mineral level.

EXAMPLE: A sodium (Na) test level of 24 mg% divided by a potassium (K) level of 10 mg% equals a Na/K ratio of 2.4 to 1.

SIGNIFICANT RATIOS

If the synergistic relationship (or ratio) between certain minerals in the body is disturbed, studies show that normal biological functions and metabolic activity can be adversely affected. Even at extremely low concentrations, the synergistic and/or antagonistic relationships between minerals still exist, which can indirectly affect metabolism.

TOXIC RATIOS

It is important to note that individuals with elevated toxic levels may not always exhibit clinical symptoms associated with those particular toxic minerals. However, research has shown that toxic minerals can also produce an antagonistic effect on various essential minerals eventually leading to disturbances in their metabolic utilization.

ADDITIONAL RATIOS

These ratios are being reported solely for the purpose of gathering research data. This information will then be used to help the attending health-care professional in evaluating their impact upon health.

REFERENCE INTERVALS

Generally, reference intervals should be considered as guidelines for comparison with the reported test values. These reference intervals have been statistically established from studying an international population of "healthy" individuals.

Important Note: The reference intervals should not be considered as absolute limits for determining deficiency, toxicity or acceptance.

THE FOLLOWING RECOMMENDATIONS SHOULD BE TAKEN ONLY WITH MEALS IN ORDER TO INCREASE ABSORPTION AND TO AVOID STOMACH DISCOMFORT. IF DISCOMFORT OCCURS SUPPLEMENTATION CAN BE REDUCED TO A MINIMUM THEN INCREASED GRADUALLY.

RECOMMENDATION	AM	NOON	PM
SYM BALANCE	1	0	1
ADEN COMPLEX	1	1	1
COPPER PLUS	1	0	1
DIGESTIVE-ZYME	1	1	1
SEL-E PLUS	1	1	1

THESE RECOMMENDATIONS ARE BASED UPON THE MINERAL LEVELS FOUND IN THE HAIR TISSUE MINERAL ANALYSIS AND MAY AT TIMES NEED MODIFICATION AS PER SPECIFIC NEED AND/OR INDIVIDUAL CIRCUMSTANCES. THESE RECOMMENDATIONS ARE PROVIDED ONLY AS A PROFESSIONAL GUIDE TO SUPPLEMENTAL ASSISTANCE.

THESE RECOMMENDATIONS MAY NOT INCLUDE MINERALS WHICH APPEAR BELOW NORMAL OR IN TURN MAY RECOMMEND MINERALS WHICH APPEAR ABOVE NORMAL ON THE HTMA GRAPH. THIS IS NOT AN OVERSIGHT. SPECIFIC MINERALS WILL INTERACT WITH OTHER MINERALS TO RAISE OR LOWER TISSUE MINERAL LEVELS, AND THIS PROGRAM IS DESIGNED TO BALANCE THE PATIENT'S MINERAL LEVELS THROUGH THESE INTERACTIONS.

THESE RECOMMENDATIONS SHOULD NOT BE TAKEN OVER A PROLONGED PERIOD OF TIME WITHOUT OBTAINING A RE-EVALUATION. THIS IS NECESSARY IN ORDER TO MONITOR PROGRESS AND MAKE THE NECESSARY CHANGES IN THE NUTRITIONAL RECOMMENDATIONS AS REQUIRED.

SPECIAL NOTE: NUTRITIONAL SUPPLEMENTS DO NOT TAKE THE PLACE OF A GOOD DIET. THEY ARE BUT AN ADDITIONAL SOURCE OF NUTRIENTS, AND THEREFORE, MUST NOT BE SUBSTITUTED FOR A BALANCED DIET.

INTRODUCTION

THE FOLLOWING REPORT SHOULD NOT BE CONSIDERED AS DIAGNOSTIC, BUT RATHER AS A SCREENING TOOL THAT PROVIDES AN ADDITIONAL SOURCE OF INFORMATION. THIS REPORT SHOULD ONLY BE USED IN CONJUNCTION WITH OTHER LABORATORY TESTS, HISTORY, PHYSICAL EXAMINATION AND THE CLINICAL EXPERTISE OF THE ATTENDING HEALTHCARE PROFESSIONAL.

TEST RESULTS WERE OBTAINED BY A LICENSED* CLINICAL LABORATORY ADHERING TO TESTING PROCEDURES THAT COMPLY WITH GOVERNMENTAL PROTOCOL AND STANDARDS ESTABLISHED BY TRACE ELEMENTS, INC., U.S.A. THE FOLLOWING INTERPRETATION IS BASED UPON INTERNATIONAL DATA AND DEFINED BY EXTENSIVE CLINICAL RESEARCH CONDUCTED BY DAVID L. WATTS, PH.D.

This analysis including levels, ratios, ranges and recommendations are based upon the sample and sampling technique meeting the following requirements:

- ** Sample obtained from the mid-parietal to the occipital region of scalp.
- ** Sample is proximal portion of hair length (first 1" to 2" of hair closest to scalp.
- ** Sufficient sample weight (minimum of 150 mg.)
- ** High grade stainless steel sampling scissors.
- ** Untreated virgin hair (no recent perms, bleaching, or coloring agents).

* Clinical Laboratory License

U.S. Department of Health and Human Services, State of Texas Department of Health,

Clinical Laboratories Improvement Act, 1988 No. 45-D0481787

METABOLIC TYPE

FAST METABOLIZER, TYPE #4

The patient is classified as a FAST METABOLIZER TYPE #4. This metabolic type has a dominance of phosphorus relative to calcium (sympathetic dominance), with an existing adrenal and thyroid insufficiency. This pattern is characteristic of "stress burnout," which can be a result of prolonged, chronic stress. This pattern may result in extreme fatigue and depression.

Endocrine replacement therapy, such as; thyroid, insulin, adrenal steroids (anti-inflammatory drugs), etc., as well as endocrine antagonists and in extreme cases of surgical removal of a gland, can affect the tissue mineral pattern. In these cases, the above reported indications of endocrine status should not be considered as representative of endocrine activity. Additional clinical tests and patient history should be taken into consideration.

There are several sub-classifications of each metabolic type, ranging from Type #1 to Type #4. This is taken into consideration on their supplement and dietary recommendations. The extent to which the patient is manifesting these metabolic characteristics depends upon the degree and chronicity of the mineral patterns.

RE-EVALUATION

A re-evaluation is suggested at two months from the beginning of implementation of the supplement program. The metabolic subtypes, such as the Type #4 may result from an acute condition, and therefore, may show a metabolic response more quickly than the Type #1.

TRENDS

The following trends may or may not be manifesting in the patient at this time. Each trend that is listed is a result of research including statistical and clinical observations. This trend analysis is advanced merely for the consideration of the health professional, and should not be considered an assessment of a medical condition. Further investigation may be indicated based upon your own clinical evaluation.

*** SPECIAL NOTE ***

It must be emphasized that the following are only trends of potential health conditions. Realistically, the probability for each trend's occurrence is based upon the degree and duration of the specific mineral imbalance. Since this analysis is not capable of determining either the previous degree of imbalance and/or previous duration, the trend analysis should only be used as an indicator to the health-care professional of potential manifestation's, particularly if the biochemical imbalance continues.

TENDENCY	1	2	3	4	5	6	7	8
ANEMIA	██████████							
ANXIETY	██████████							
FATIGUE	██████████							
INSOMNIA	██							

COMMENTS

ANEMIA AND LOW IRON:

The patient's tissue iron level is low. This is indicative of a trend toward iron deficiency anemia. Serum changes that may be evaluated are:

Hypochromic Anemia
Low Hematocrit

Microcytic Anemia
Low Hemoglobin

These serum indications may be present only if iron deficiency is chronic. If iron intake is marginally adequate, the above indications may not be manifesting, but iron deficiency symptoms may be present. Iron deficiency anemia symptoms prior to iron deficiency are not uncommon. If the patient's plasma transferrin is less than 10% saturated, it can be an early indicator of an eventual iron deficiency anemia, even if all other parameters are normal.

ANXIETY:

Low tissue calcium is associated with increased central nervous system sensitivity and increased serum lactic acid levels, both of which may contribute to increased anxiety states. Anxiety may be contributed to by any factor that interferes with normal calcium metabolism such as stress or accumulation of toxic metals such as lead and mercury.

FATIGUE AND LOW IRON:

Low tissue iron is associated with a tendency toward anemia. Iron deficiency anemia is a contributing factor to fatigue and shortness of breath.

INSOMNIA:

Two types of insomnia should be distinguished in order to determine effective treatment.

INSOMNIA AND CALCIUM:

Insomnia characterized by an inability to fall asleep is most often associated with an increased need for calcium. If calcium is not recommended on the patient's program, calcium may be given until symptoms have improved.

CONTRAINDICATIONS

It is suggested that additional supplementation and/or intake of the following nutrients and food substitutes (if any) should be avoided by the patient until re-evaluation.

*** ASPARTAME ***

Aspartame (aspartic acid) is an ingredient of some sweeteners. In the metabolic pathways, aspartic acid is converted to phenylalanine, which is further metabolized to tyrosine. In the presence of an iron deficiency, phenylalanine is not metabolized to tyrosine. The amino acid phenylalanine has a sedative effect that, if excessive, may contribute to fatigue and depression.

DIETARY SUGGESTIONS

The following dietary suggestions are defined by several factors: the individual's mineral levels, ratios and metabolic type, as well as the nutrient value of each food including protein, carbohydrate, fat, and vitamin and mineral content. Based upon these determinations, it may be suggested that foods be avoided or increased temporarily in the diet to aid in the improvement of the patient's chemistry.

GENERAL DIETARY GUIDELINES FOR THE FAST METABOLIZER

* INCREASE INTAKE OF HIGH PURINE PROTEIN FOODS...high purine protein sources include liver, kidney and heart. Other good sources include sardines, tuna, clams, crab, lobster and oysters. Unless notified otherwise, high purine and moderate purine protein intake should constitute approximately 33% of total daily caloric intake.

* INCREASE INTAKE OF MILK AND MILK PRODUCTS...such as cheese, yogurt, cream, butter (unsalted). Increase intake of nuts and seeds such as almonds, walnuts, peanuts, peanut butter and sunflower seeds. Foods high in fat unless notified otherwise should constitute approximately 33% of total daily caloric intake.

* REDUCE CARBOHYDRATE INTAKE...including unrefined carbohydrates. Sources such as cereals, whole grains and whole grain products are contraindicated for frequent consumption until the next evaluation. Carbohydrate intake in the form of unrefined carbohydrates should be approximately 33% of total daily caloric intake.

* AVOID ALL SUGARS AND REFINED CARBOHYDRATES...this includes white and brown sugar, honey, candy, soda pop, cake, pastries, alcohol and white bread.

FOOD ALLERGIES:

In some individuals, certain foods can produce a maladaptive or "allergic-like" reaction commonly called "food allergies". Consumption of foods that one is sensitive to can bring about reactions ranging from drowsiness to hyperactivity in children, itching and rashes, headaches, high-blood pressure and arthritic pain.

Sensitivity to foods can develop due to biochemical (nutritional) imbalances, and which stress, pollution, and medications can aggravate. Nutritional imbalance can further be contributed to by restricting food variety, such as eating only a small group of foods on a daily basis. Often a person will develop a craving for the food they are most sensitive to and may eat the same food or food group more than once a day.

The following section may contain foods that are recommended to avoid. These foods should be considered as potential "allergy foods", or as foods that may impede a rapid and effective response. Consumption of these foods should be avoided completely for four days. Afterwhich, they should not be eaten more frequently than once every three days during course of therapy.

FOODS THAT STIMULATE HISTAMINES:

Consumption of the following foods can stimulate histamine release in certain metabolic types and may contribute to respiratory-type allergy reactions. These foods are to be avoided until the next evaluation or until notified otherwise by attending doctor.

Beet Greens	Rhubarb
Apples	Chocolate
Spinach	Black Tea
Eggplant	Strawberries
Sweet Potatoes	Peanuts
Blueberries	Green Beans

Pecans
Wheat Germ
Cocoa
Parsley
Beets

Chard
Concord Grapes
Collards
Blackberries

THE FOLLOWING FOODS MAY BE INCREASED IN THE DIET UNTIL THE NEXT EVALUATION:

Mozzarella Cheese
Milk
Kale
Monterey Cheese
Almonds
Swiss Cheese

Turnip Greens
Mustard Greens
Yogurt
Cream
Buttermilk

FOODS HIGH IN IRON CONTENT:

The following foods may be increased in the diet:

Beef (lean)
Egg (yolk)
Pumpkin Seeds

Sunflower Seeds
Oysters
Chipped Beef

NOTE: Dairy foods such as milk and cheese actually decrease the availability of iron from the diet. If dairy foods or vegetarian diets are consumed exclusive of meats, iron absorption can be reduced by as much as 60 percent.

AMINO ACIDS THAT IMPROVE CALCIUM ABSORPTION:

Calcium absorption is greatly enhanced when the diet is high in the amino acids, lysine, arginine and histadine. These proteins also help to reduce acidity of the tissues. Both effects are favorable for the fast metabolizer, therefore addition of any of the following foods to the diet is recommended at this time:

Lima Beans
Garbanzo Beans
Rumproast
Skim Milk
Beef Stew
Cottage Cheese
Spare Ribs
Lentils
Flounder
Cod
Ham

Salami
Sausage (lean)
Lamb
Smelt
Vegetable Stew
Canadian bacon
Peanuts
Bass
Heart
Chuck Roast
Liverwurst

SPECIAL NOTE:

This analysis will list only a limited number of dietary foods to avoid or to increase in the diet. For those foods not specifically mentioned in this section, continued consumption on a moderate basis may be considered appropriate unless recommended otherwise.

NO PART OF THIS INTERPRETIVE REPORT MAY BE REPRODUCED OR TRANSMITTED IN ANY FORM OR BY ANY MEANS, ELECTRONIC OR MECHANICAL, INCLUDING PHOTOCOPYING, RECORDING, OR ANY INFORMATION STORAGE OR RETRIEVAL SYSTEM WITHOUT PERMISSION IN WRITING FROM TRACE ELEMENTS, INC., U.S.A.

InterClinical Laboratories Pty Limited
Unit 6/10 Bradford Street, Alexandria, N.S.W. 2015, Sydney, Australia
Ph: (02) 9693-2888 Fax: (02) 9693-1888
Email: lab@interclinical.com.au

Authorized Representative for Australia and New Zealand